



# SAMHSA's Concept of Trauma and Guidance for a Trauma-Informed Approach

Prepared by  
SAMHSA's Trauma and Justice Strategic Initiative  
July 2014



## Introduction

Trauma is a widespread, harmful and costly public health problem. It occurs as a result of violence, abuse, neglect, loss, disaster, war and other emotionally harmful experiences. Trauma has no boundaries with regard to age, gender, socioeconomic status, race, ethnicity, geography or sexual orientation. It is an almost universal experience of people with mental and substance use disorders. The need to address trauma is increasingly viewed as an important component of effective behavioral health service delivery. Additionally, it has become evident that addressing trauma requires a multi-pronged, multi-agency public health approach inclusive of public education and awareness, prevention and early identification, and effective trauma-specific assessment and treatment. **In order to maximize the impact of these efforts, they need to be provided in an organizational or community context that is trauma-informed, that is, based on the knowledge and understanding of trauma and its far-reaching implications.**

**The need to address trauma is increasingly viewed as an important component of effective behavioral health service delivery.**

The effects of traumatic events place a heavy burden on individuals, families and communities and create challenges for public institutions and service systems. Although many people who experience a traumatic event will go on with their lives without lasting negative effects, others will have more difficulty and experience traumatic stress reactions. Emerging research has documented the relationships among exposure to traumatic events, impaired neurodevelopmental and immune systems responses and subsequent health risk behaviors resulting in chronic physical and behavioral health disorders.<sup>1,2,3,4,5</sup> Research has also indicated that with appropriate

support and intervention, people can overcome traumatic experiences.<sup>6,7,8,9</sup> However, most people go without these services and supports. Unaddressed trauma significantly increases the risk of mental and substance use disorders and chronic physical diseases.<sup>1,10,11</sup>

**With appropriate supports and intervention, people can overcome traumatic experiences.**

Individuals with experiences of trauma are found in multiple service sectors, not just in behavioral health. Studies of people in the juvenile and criminal justice system reveal high rates of mental and substance use disorders and personal histories of trauma.<sup>12,13</sup> Children and families in the child welfare system similarly experience high rates of trauma and associated behavioral health problems.<sup>14</sup> Young people bring their experiences of trauma into the school systems, often interfering with their school success. And many patients in primary care similarly have significant trauma histories which has an impact on their health and their responsiveness to health interventions.<sup>15,16,17</sup>

In addition, the public institutions and service systems that are intended to provide services and supports to individuals are often themselves trauma-inducing. The use of coercive practices, such as seclusion and restraints, in the behavioral health system; the abrupt removal of a child from an abusing family in the child welfare system; the use of invasive procedures in the medical system; the harsh disciplinary practices in educational/school systems; or intimidating practices in the criminal justice system can be re-traumatizing for individuals who already enter these systems with significant histories of trauma. These program or system practices and policies often interfere with achieving the desired outcomes in these systems.

**There is an increasing focus on the impact of trauma and how service systems may help to resolve or exacerbate trauma-related issues. These systems are beginning to revisit how they conduct their business under the framework of a trauma-informed approach.**

page 2

Thus, the pervasive and harmful impact of traumatic events on individuals, families and communities and the unintended but similarly widespread re-traumatizing of individuals within our public institutions and service systems, makes it necessary to rethink doing "business as usual." In public institutions and service systems, there is increasing recognition that many of the individuals have extensive histories of trauma that, left unaddressed, can get in the way of achieving good health and well-being. For example, a child who suffers from maltreatment or neglect in the home may not be able to concentrate on school work and be successful in school; a woman victimized by domestic violence may have trouble performing in the work setting; a jail inmate repeatedly exposed to violence on the street may have difficulty refraining from retaliatory violence and re-offending; a sexually abused homeless youth may engage in self-injury and high risk behaviors to cope with the effects of sexual abuse; and, a veteran may use substances to mask the traumatic memories of combat. The experiences of these individuals are compelling and, unfortunately, all too common. Yet, until recently, gaining a better understanding of how to address the trauma

experienced by these individuals and how to mitigate the re-traumatizing effect of many of our public institutions and service settings was not an integral part of the work of these systems. Now, however, there is an increasing focus on the impact of trauma and how service systems may help to resolve or exacerbate trauma-related issues. These systems are beginning to revisit how they conduct their "business" under the framework of a trauma-informed approach.

## Purpose and Approach: Developing a Framework for Trauma and a Trauma-Informed Approach

**PURPOSE**  
The purpose of this paper is to develop a working concept of trauma and a trauma-informed approach and to develop a shared understanding of these concepts that would be acceptable and appropriate across an array of service systems and stakeholder groups. SAMHSA puts forth a framework for the behavioral health specialty sectors, that can be adapted to other sectors such as child welfare, education, criminal and juvenile justice, primary health care, the military and other settings that have the potential to ease or exacerbate an individual's capacity to cope with traumatic experiences. In fact, many people with behavioral health problems receive treatment and services in these non-specialty behavioral health systems. SAMHSA intends this

framework to be relevant to its federal partners and their state and local system counterparts and to practitioners, researchers, and trauma survivors, families and communities. The desired goal is to build a framework that helps systems "talk" to each other, to understand better the connections between trauma and behavioral health issues, and to guide systems to become trauma-informed.

**APPROACH**  
SAMHSA approached this task by integrating three significant threads of work: trauma focused research work; practice-generated knowledge about trauma interventions; and the lessons articulated by survivors

**There is an increasing focus on the impact of trauma and how service systems may help to resolve or exacerbate trauma-related issues. These systems are beginning to revisit how they conduct their business under the framework of a trauma-informed approach.**

page 3

of traumatic experiences who have had involvement in multiple service sectors. It was expected that this blending of the research, practice and survivor knowledge would generate a framework for improving the capacity of our service systems and public institutions to better address the trauma-related issues of their constituents.

To begin this work, SAMHSA conducted an environmental scan of trauma definitions and models of trauma informed care. SAMHSA convened a group of national experts who had done extensive work in this area. This included trauma survivors who had been recipients of care in multiple service systems; practitioners from an array of fields, who had experience in trauma treatment; researchers whose work focused on trauma and the development of trauma-specific interventions; and policymakers in the field of behavioral health.

From this meeting, SAMHSA developed a working document summarizing the discussions among these experts. The document was then vetted among federal agencies that conduct work in the field of trauma. Simultaneously, it was placed on a SAMHSA website for public comment. Federal agency experts provided rich comments and suggestions; the public comment site drew just over 2,000 respondents and 20,000 comments or endorsements of others' comments. SAMHSA reviewed all of these comments, made revisions to the document and developed the framework and guidance presented in this paper.

SAMHSA's approach to this task has been an attempt to integrate knowledge developed through research and clinical practice with the voices of trauma survivors. This also included experts funded through SAMHSA's trauma-focused grants and initiatives, such as SAMHSA's National Child Traumatic Stress Initiative, SAMHSA's National Center for Trauma Informed Care, and data and lessons learned from other grant programs that did not have a primary focus on trauma but included significant attention to trauma, such as SAMHSA's Jail Diversion Trauma Recovery grant program; Children's Mental Health Initiative; Women, Children and Family Substance Abuse Treatment Program; and Offender Reentry and Adult Treatment Drug Court Programs.

**The key questions addressed in this paper are:**

- What do we mean by trauma?
- What do we mean by a trauma-informed approach?
- What are the key principles of a trauma-informed approach?
- What is the suggested guidance for implementing a trauma-informed approach?
- How do we understand trauma in the context of community?

of traumatic experiences who have had involvement in multiple service sectors. It was expected that this blending of the research, practice and survivor knowledge would generate a framework for improving the capacity of our service systems and public institutions to better address the trauma-related issues of their constituents.

To begin this work, SAMHSA conducted an environmental scan of trauma definitions and models of trauma informed care. SAMHSA convened a group of national experts who had done extensive work in this area. This included trauma survivors who had been recipients of care in multiple service systems; practitioners from an array of fields, who had experience in trauma treatment; researchers whose work focused on trauma and the development of trauma-specific interventions; and policymakers in the field of behavioral health.

From this meeting, SAMHSA developed a working document summarizing the discussions among these experts. The document was then vetted among federal agencies that conduct work in the field of trauma. Simultaneously, it was placed on a SAMHSA website for public comment. Federal agency experts provided rich comments and suggestions; the public comment site drew just over 2,000 respondents and 20,000 comments or endorsements of others' comments. SAMHSA reviewed all of these comments, made revisions to the document and developed the framework and guidance presented in this paper.

SAMHSA's approach to this task has been an attempt to integrate knowledge developed through research and clinical practice with the voices of trauma survivors. This also included experts funded through SAMHSA's trauma-focused grants and initiatives, such as SAMHSA's National Child Traumatic Stress Initiative, SAMHSA's National Center for Trauma Informed Care, and data and lessons learned from other grant programs that did not have a primary focus on trauma but included significant attention to trauma, such as SAMHSA's Jail Diversion Trauma Recovery grant program; Children's Mental Health Initiative; Women, Children and Family Substance Abuse Treatment Program; and Offender Reentry and Adult Treatment Drug Court Programs.

## Background: Trauma — Where We Are and How We Got Here

The concept of traumatic stress emerged in the field of mental health at least four decades ago. Over the last 20 years, SAMHSA has been a leader in recognizing the need to address trauma as a fundamental obligation for public mental health and substance abuse service delivery and has supported the development and promulgation of trauma-informed systems of care. In 1994, SAMHSA convened the Baring to Vision Conference, an event designed to bring trauma to the foreground and the first national conference in which women trauma survivors talked about their experiences and ways in which standard practices in hospitals re-traumatized and often, triggered memories of previous abuse. In 1998, SAMHSA funded the Women, Co-Occurring Disorders and Violence Study to generate knowledge on the development and evaluation of integrating services and approaches for women with co-occurring mental and substance use disorders who also had histories of physical and/or sexual abuse. In 2001, SAMHSA funded the National Child Traumatic Stress Initiative to increase understanding of child trauma and develop effective interventions for children exposed to different types of traumatic events.

The American Psychiatric Association (APA) created an important role in defining trauma. Diagnostic criteria for traumatic stress disorders have been debated through several iterations of the Diagnostic and Statistical Manual of Mental Disorders (DSM) with a new category of Trauma- and Stressor-Related Disorders, across the life-span, included in the recently released DSM-V (APA, 2013). Measures and inventories of trauma exposure, with both clinical and research applications, have proliferated since the 1970s.<sup>18,19,20,21</sup> National trauma research and practice centers have conducted significant work in the past few decades, further refining the concept of trauma, and developing effective trauma assessments and treatments.<sup>22,23,24,25</sup> With the advances in neuroscience, a biopsychosocial approach to traumatic experiences has begun to delineate the mechanisms in which neurobiology, psychological processes, and social attachment interact and contribute to mental and substance use disorders across the life-span.<sup>3,25</sup>

Simultaneously, an emerging trauma survivors movement has provided another perspective on the understanding of traumatic experiences. Trauma survivors, that is, people with lived experience of trauma, have powerfully and systematically documented their paths to recovery.<sup>26</sup> Traumatic experiences complicate a child's or an adult's capacity to make sense of their lives and to create meaningful consistent relationships in their families and communities.

**Trauma survivors have powerfully and systematically documented their paths to recovery.**

With the growing understanding of the pervasiveness of traumatic experience and responses, a growing number of clinical interventions for trauma responses have been developed. Federal research agencies, academic institutions and practice-research partnerships have generated empirically-supported interventions. In SAMHSA's National Registry of Evidence-based Programs and Practices (NREPP) alone there are over 15 interventions focusing on the treatment or recovery for trauma.

These interventions have been integrated into the behavioral health treatment care delivery system; however, from the voice of trauma survivors, it has become clear that these clinical interventions are not enough. Building on lessons learned from SAMHSA's Women, Co-Occurring Disorders and Violence Study; SAMHSA's National Child Traumatic Stress Network; and SAMHSA's National Center for Trauma-Informed Care and Alternatives to Seclusion and Restraints, among other developments in the field, it became clear that the organizational climate and conditions in which services are provided played a significant role in maximizing the outcomes of interventions and contributing to the healing and recovery of the people being served. SAMHSA's National Center for Trauma-Informed Care has continued to advance this effort, starting first in the behavioral health sector, but increasingly responding to technical assistance requests for organizational change in the criminal justice, education, and primary care sectors.

The convergence of the trauma survivor's perspective with research and clinical work has underscored the central role of traumatic experiences in the lives of people with mental and substance use conditions. The connection between trauma and these conditions offers a potential explanatory model for what has happened to individuals, both children and adults, who come to the attention of the behavioral health and other service systems.<sup>25,27</sup>

People with traumatic experiences, however, do not show up only in behavioral health systems. Responses to these experiences often manifest in behaviors or conditions that result in involvement with the child welfare and the criminal and juvenile justice system or in difficulties in the education, employment or primary care system. Recently, there has also been a focus on individuals in the military and increasing rates of posttraumatic stress disorders.<sup>28,29,30,31</sup>

**FEDERAL, STATE AND LOCAL LEVEL TRAUMA-FOCUSED ACTIVITIES**  
The increased understanding of the pervasiveness of trauma and its connections to physical and behavioral health and well-being, have propelled a growing number of organizations and service systems to explore ways to make their services more responsive to people who have experienced trauma. This has been happening in state and local systems and federal agencies.

States are elevating a focus on trauma. For example, Oregon Health Authority is looking at different types of trauma across the age span and different population groups. Maine's "Thrive Initiative" incorporates a

trauma-informed care focus in their children's systems of care. New York is introducing a trauma-informed element in the juvenile justice system. Missouri is exploring a trauma-informed approach for their adult mental health system. In Massachusetts, the Child Trauma Project is focused on taking trauma-informed care statewide in child welfare practice. In Connecticut the Child Health and Development Institute with the state Department of Children and Families is building a trauma-informed system of care throughout the state through policy and workforce development. SAMHSA has supported the further development of trauma-informed approaches through its Mental Health Transformation Grant program directed to State and local governments.

Increasing examples of local level efforts are being documented. For example, the City of Barton Springs in Florida has taken significant steps in becoming a trauma-informed community. The agency made it its mission to promote a widespread awareness of the costly effects of personal adversity upon the wellbeing of the community. The Family Policy Council in Washington State convened groups to focus on the impact of adverse childhood experiences on the health and well-being of its local communities and tribal communities. Philadelphia held a summit to further its understanding of the impact of trauma and violence on the psychological and physical health of its communities.

**SAMHSA continues its support of grant programs that specifically address trauma.**

At the federal level, SAMHSA continues its support of grant programs that specifically address trauma and technical assistance centers that focus on prevention, treatment and recovery from trauma.

Other federal agencies have increased their focus on trauma. The Administration on Children Youth and Families (ACYF) has focused on the complex trauma of children in the child welfare system and how screening and assessing for severity of trauma and linkage with trauma treatments can contribute to improved well-being for these youth. In a joint effort among ACYF, SAMHSA and the Centers for Medicare and Medicaid Services (CMS), the three agencies developed and issued through the CMS State Directors' mechanism, a letter to all State Child Welfare Administrators, Mental Health Commissioners, Single State Agency Directors for Substance Abuse and State Medicaid Directors discussing trauma, its impact on children, screening, assessment and treatment interventions and strategies for paying for such care. The Office of Juvenile Justice and Delinquency Prevention has specific recommendations to address trauma in their Children Exposed to Violence Initiative. The Office of Women's Health has developed a curriculum to train providers in

health care on how to address trauma issues in health care for women. The Department of Labor is examining trauma and the workplace through a federal interagency workgroup. The Department of Defense is honing in on prevention of sexual violence and trauma in the military.

As multiple federal agencies recognized varied sectors have recognized the impact of traumatic experiences on the children, adults, and families they serve, they have requested collaboration with SAMHSA in addressing these issues. The widespread recognition of the impact of trauma and the burgeoning interest in developing capacity to respond through trauma-informed approaches compelled SAMHSA to revisit its conceptual framework and approach to trauma, as well as its applicability not only to behavioral health but also to other related fields.

On a national level, SAMHSA continues its support of grant programs that specifically address trauma and technical assistance centers that focus on prevention, treatment and recovery from trauma.

States are elevating a focus on trauma. For example, Oregon Health Authority is looking at different types of trauma across the age span and different population groups. Maine's "Thrive Initiative" incorporates a

trauma-informed care focus in their children's systems of care. New York is introducing a trauma-informed element in the juvenile justice system. Missouri is exploring a trauma-informed approach for their adult mental health system. In Massachusetts, the Child Trauma Project is focused on taking trauma-informed care statewide in child welfare practice. In Connecticut the Child Health and Development Institute with the state Department of Children and Families is building a trauma-informed system of care throughout the state through policy and workforce development. SAMHSA has supported the further development of trauma-informed approaches through its Mental Health Transformation Grant program directed to State and local governments.

Increasing examples of local level efforts are being documented. For example, the City of Barton Springs in Florida has taken significant steps in becoming a trauma-informed community. The agency made it its mission to promote a widespread awareness of the costly effects of personal adversity upon the wellbeing of the community. The Family Policy Council in Washington State convened groups to focus on the impact of adverse childhood experiences on the health and well-being of its local communities and tribal communities. Philadelphia held a summit to further its understanding of the impact of trauma and violence on the psychological and physical health of its communities.

At the federal level, SAMHSA continues its support of grant programs that specifically address trauma and technical assistance centers that focus on prevention, treatment and recovery from trauma.

States are elevating a focus on trauma. For example, Oregon Health Authority is looking at different types of trauma across the age span and different population groups. Maine's "Thrive Initiative" incorporates a

trauma-informed care focus in their children's systems of care. New York is introducing a trauma-informed element in the juvenile justice system. Missouri is exploring a trauma-informed approach for their adult mental health system. In Massachusetts, the Child Trauma Project is focused on taking trauma-informed care statewide in child welfare practice. In Connecticut the Child Health and Development Institute with the state Department of Children and Families is building a trauma-informed system of care throughout the state through policy and workforce development. SAMHSA has supported the further development of trauma-informed approaches through its Mental Health Transformation Grant program directed to State and local governments.

page 4

page 5

page 6

page 7

## Endnotes

- <sup>1</sup> Felitti, G., Anda, R., Nordenberg, D., et al., (1998). Relationship of child abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences Study. *American Journal of Preventive Medicine*, 14, 245-258.
- <sup>2</sup> Anda, R.F., Brown, D.W., Dube, S.R., Bremner, J.D., Felitti, V.J., and Giles, W.G. (2008). Adverse childhood experiences and chronic obstructive pulmonary disease in adults. *American Journal of Preventive Medicine*, 34(5), 396-403.
- <sup>3</sup> Perry, B., (2004). Understanding traumatized and maltreated children: The core concepts – Living and working with traumatized children. The Child Trauma Academy. [www.ChildTrauma.org](http://www.ChildTrauma.org).
- <sup>4</sup> Shonkoff, J.P., Garner, A.S., Siegel, B.S., Dobbins, M.L., Earls, M.F., McGuinn, L., et al., Wood, D.L. (2012). The lifelong effects of early childhood adversity and toxic stress. *Pediatrics*, 129(1), 232-246.
- <sup>5</sup> McLoughlin, K.A., Green, J.G., Kessler, R.C., et al. (2009). Childhood adversity and adult psychiatric disorder in the US National Comorbidity Survey. *Psychol Med*, 40(4), 847-59.
- <sup>6</sup> National Child Traumatic Stress Network Systems Integration Working Group (2005). Helping children in the child welfare system heal from trauma: A systems integration approach.
- <sup>7</sup> Dozier, M., Cue, K.L., and Barnett, L. (1994). Clinicians as caregivers: Role of attachment organization in treatment. *Journal of Consulting and Clinical Psychology*, 62(4), 793-800.
- <sup>8</sup> Najavits, L.M. (2002). *Seeking Safety: A Treatment Manual for PTSD and Substance Abuse*. New York: Guilford Press.
- <sup>9</sup> Covington, S. (2008). "Women and Addiction: A Trauma-Informed Approach." *Journal of Psychoactive Drugs*. SARC Supplement 5, November 2008, 377-385.
- <sup>10</sup> Anda, R.F., Brown, D.W., Dube, S.R., Bremner, J.D., Felitti, V.J., and Giles, W.H. (2008). Adverse childhood experiences and chronic obstructive pulmonary disease in adults. *American Journal of Preventive Medicine*, 34(5), 396-403.
- <sup>11</sup> Dube, S.R., Felitti, V.J., Dong, M., Chapman, D.P., Giles, W.H., and Anda, R.F. (2003). Childhood abuse, neglect, and household dysfunction and the risk of illicit drug use: The Adverse Childhood Experiences Study. *Pediatrics*, 111(3), 564-572.
- <sup>12</sup> Ford, J. and Wilson, C. (2012). SAMHSA's Trauma and Trauma-Informed Care Experts Meeting.
- <sup>13</sup> Ford, J.D. (2013). Treatment of complex trauma: A sequenced, relationship-based approach. New York, NY, US: Guilford Press.
- <sup>14</sup> Wilson, C., and Conradi, L. (2010). Managing traumatized children: A trauma systems perspective. *Psychiatry*. doi:10.1097/MOP.0b013e31832833e0766
- <sup>15</sup> Dutton, M.A., Bonnie, L.G., Kaltman, S.I., Roesch, D.M., and Zeffiro, T.A., et al. (2006). Intimate partner violence, PTSD, and adverse health outcomes. *Journal of Interpersonal Violence*, 21(7), 955-968.
- <sup>16</sup> Campbell, R., Greeson, M.R., Bybee, D., and Raja, S. (2008). The co-occurrence of childhood sexual abuse, adult sexual assault, intimate partner violence, and sexual harassment: A mediational model of posttraumatic stress disorder and physical health outcomes. *Journal of Consulting and Clinical Psychology*, 76(2), 194-207.
- <sup>17</sup> Bonomi, A.E., Anderson, M.L., Rivara, F.P., Thompson, R.S. (2007). Health outcomes in women with physical and sexual intimate partner violence exposure. *Journal of Women's Health*, 16(7), 987-997.
- <sup>18</sup> Norris, F.H. (1990). Screening for traumatic stress: A scale for use in the general population. *Journal of Applied Social Psychology*, 20, 1704-1718.
- <sup>19</sup> Norris, F.H. and Hamblen, J.L. (2004). Standardized self-report measures of civilian trauma and PTSD. In J.P. Wilson, T.M. Keane and T. Martin (Eds.), *Assessing psychological trauma and PTSD* (pp. 63-102). New York: Guilford Press.
- <sup>20</sup> Orsillo, S.M. (2001). Measures for acute stress disorder and posttraumatic stress disorder. In M.M. Antony and S.M. Orsillo (Eds.), *Practitioner's Guide to Empirically Based Measures of Anxiety* (pp. 255-307). New York: Kluwer Academic/Plenum
- <sup>21</sup> Weathers, F.W. and Keane, T.M. (2007). The criterion A problem revisited: Controversies and challenges in defining and measuring psychological trauma. *Journal of Traumatic Stress*, 20(2), 107-121.
- <sup>22</sup> Van der Kolk, B. (2003). The neurobiology of childhood trauma and abuse. Laor, N. and Wolmer, L. (guest editors): *Child and Adolescent Psychiatric Clinics of North America: Posttraumatic Stress Disorder*, 12 (2), Philadelphia: W.B. Saunders, 293-317.
- <sup>23</sup> Herman, J. (1992). *Trauma and recovery: The aftermath of violence – from domestic abuse to political terror*. New York: Basic Books.
- <sup>24</sup> Harris, M., and Fallot, R. (2001). Using trauma theory to design service systems. *New Directions for Mental Health Services*, 89, Jossey Bass.
- <sup>25</sup> Bloom, S. (2012). "The Workplace and trauma-informed systems of care." Presentation at the National Network to Eliminate Disparities in Behavioral Health. Cohen, J., Mannarino, A., Deblinger, E., (2004). Trauma-focused Cognitive Behavioral Therapy (TF-CBT). Available from: <http://tfcbt.musc.edu>.
- <sup>26</sup> SAMHSA's National Center for Trauma-Informed Care (2012). *Report of Project Activities Over the Past 18 Months, History, and Selected Products*. Available from: [http://www.nasmhpd.org/docs/NCTIC/NCTIC\\_Final\\_Report\\_3-26-12.pdf](http://www.nasmhpd.org/docs/NCTIC/NCTIC_Final_Report_3-26-12.pdf)
- <sup>27</sup> Bloom, S.L., and Farragher, B. (2011). *Destroying Sanctuary: The crisis in human services delivery systems*. New York: Oxford University Press. Guarino, K., Soares, P., Konath, K., Clervil, R., and Bassuk, E. (2009). *Trauma-Informed Organizational Toolkit*. Rockville, MD: Center for Mental Health Services, Substance Abuse and Mental Health Services Administration, and the Daniels Fund, the National Child Traumatic Stress Network and the W.K. Kellogg Foundation.
- <sup>28</sup> Dekel, S., Ein-Dor, T., and Zahava, S. (2012). Posttraumatic growth and posttraumatic distress: A longitudinal study. *Psychological Trauma: Theory, Research, Practice, and Policy*, 4(1), 94-101.
- <sup>29</sup> Jakupcak, M., Tull, M.T., McDermott, M.J., Kayser, D., Hunt, S., and Simpson, T. (2010). PTSD symptom clusters in relationship to alcohol misuse among Iraq and Afghanistan war veterans seeking post-deployment VA health care. *Addictive Behaviors* 35(9), 840-843.
- <sup>30</sup> Goodwin, L. and Rona, R.J. (2013) PTSD in the armed forces: What have we learned from the recent cohort studies of Iraq/Afghanistan? *Journal of Mental Health* 22(5), 397-401.
- <sup>31</sup> Wolf, E.J., Mitchell, K.S., Koenen, C.K., and Miller, M.W. (2013). Combat exposure severity as a moderator of genetic and environmental liability to post-traumatic stress disorder. *Psychological Medicine*.
- <sup>32</sup> National Analytic Center-Statistical Support Services (2012). Trauma-Informed Care *White Paper*, prepared for the Substance Abuse and Mental Health Services Administration, Center for Behavioral Health Statistics and Quality.
- <sup>33</sup> Ford, J.D., Fallot, R., and Harris, M. (2009). Group Therapy. In C.A. Courtois and J.D. Ford (Eds.), *Treating complex traumatic stress disorders: An evidence-based guide* (pp.415-440). New York, NY, US: Guilford Press.
- <sup>34</sup> Brave Heart, M.Y.H., Chase, J., Elkings, J., and Altschul, D.B. (2011). Historical trauma among indigenous peoples of the Americas: Concepts, research, and clinical considerations. *Journal of Psychoactive Drugs*, 43(4), 282-290.
- <sup>35</sup> Brown, S.M., Baker, C.N., and Wilcox, P. (2012). Risking connection trauma training: A pathway toward trauma-informed care in child congregate care settings. *Psychological Trauma: Theory, Research, Practice, and Policy*, 4(5), 507-515.
- <sup>36</sup> Farragher, B. and Yanosy, S. (2005). Creating a trauma-sensitive culture in residential treatment. *Therapeutic Communities*, 26(1), 93-109.
- <sup>37</sup> Elliott, D.E., Bjelajac, P., Fallot, R.D., Markoff, L.S., and Reed, B.G. (2005). Trauma-informed or trauma-denied: Principles and implementation of trauma-informed services for women. *Journal of Community Psychology*, 33(4), 461-477.
- <sup>38</sup> Huang, L.N., Pau, T., Flatow, R., DeVoursney, D., Afayee, S., and Nugent, A. (2012). Trauma-Informed Care Models Compendium.
- <sup>39</sup> Fallot, R. and Harris, M. (2006). *Trauma-Informed Services: A Self-Assessment and Planning Protocol*. Community Connections.
- <sup>40</sup> Henry, Black-Pond, Richardson and Vandercort. (2010). Western Michigan University, Southwest Michigan Children's Trauma Assessment Center (CTAC).
- <sup>41</sup> Hummer, V. and Dollard, N. (2010). *Creating Trauma-Informed Care Environments: An Organizational Self-Assessment*. (part of *Creating Trauma-Informed Care Environments Curriculum*) Tampa FL: University of South Florida. The Department of Child and Family Studies within the College of Behavioral and Community Sciences.
- <sup>42</sup> Penney, D. and Cave, C. (2012). *Becoming a Trauma-Informed Peer-Run Organization: A Self-Reflection Tool* (2013). Adapted for Mental Health Empowerment Project, Inc. from *Creating Accessible, Culturally Relevant, Domestic Violence- and Trauma-Informed Agencies*, ASRI and National Center on Domestic Violence, Trauma and Mental Health.